

Comparison of Age and Growth of Shovelnose Sturgeon in the Missouri and Yellowstone Rivers

SCOTT R. EVERETT*¹ AND DENNIS L. SCARNECCHIA

*Department of Fish and Wildlife Resources,
University of Idaho, Post Office Box 441136,
Moscow, Idaho 83844-1136, USA*

GREG J. POWER

*North Dakota Game and Fish Department,
100 North Bismarck Expressway, Bismarck, North Dakota 58501, USA*

CHRISTOPHER J. WILLIAMS

*Division of Statistics, University of Idaho,
Post Office Box 441104, Moscow, Idaho 83844-1104, USA*

Abstract.—During the spring and summer 1991–1995, sampling was conducted on the Missouri




FIGURE 1.—Map of the lower Yellowstone River and Lake Sakakawea, Montana and North Dakota.

habitat, alterations of the natural hydrograph, altered river temperatures, and decreases in the river's turbidity. Many remaining riverine reaches of the Missouri River now consist of a single, narrow, deep, swift-flowing channel (Hesse et al. 1989; Hesse and Mestl 1993). Reductions in the variability of discharge and the overall peak discharge have eliminated the flood pulse of spring and early summer (Hesse and Mestl 1993; Galat and Rasmussen 1995). Moreover, each reservoir serves as a sediment sink, limiting the downstream movement of the river's suspended load (Hesse and Mestl 1993) and reducing the river's turbidity (Hesse 1987). The dams themselves restrict fish movement, creating fragmented and disjunct populations.

These habitat changes have had substantial effects on much of the Missouri River fish community, especially fishes such as shovelnose sturgeon that inhabit primarily main-channel habitats (Curtis et al. 1997). Many of the specific effects are poorly understood but manifested as the decline or disappearance of a species from a river segment.

Information on age and growth of a long-lived species, such as shovelnose sturgeon, can reveal the effect of habitat changes over decades. The objective of this paper is to compare growth rates of shovelnose sturgeon inhabiting two distinct segments of the Missouri River Basin—a slightly altered habitat and a highly altered habitat.

Study Area

The study area included the Yellowstone River near Intake, Montana, and in the Missouri River near Bismarck, North Dakota (Figure 1). The Yellowstone and Missouri rivers are particularly useful for comparing growth rates of shovelnose sturgeon between quasi-natural and highly altered river segments. The lower Yellowstone River has no

large main-stem dams, a near natural hydrograph, high turbidity, and habitat conditions characteristic of a slightly altered large river system (White and Bramblett 1993). The Missouri River has a similar habitat from its confluence with the Yellowstone River downstream to the headwaters of Lake Sakakawea. This entire area (hereafter called the Williston–Yellowstone segment) is inhabited by sturgeon that are able to move freely within the area. In contrast, the Missouri River in North Dakota downstream of Garrison Dam (hereafter called the Bismarck segment) is characterized as highly altered and has a highly regulated hydrograph, reduced turbidity, and cool water from hypolimnetic releases. These changes have occurred since the dam was closed in December 1953. The Bismarck segment has also undergone more bank stabilization and other habitat modifications than the Williston–Yellowstone segment. Garrison Dam effectively isolates the two populations of sturgeon. Habitat conditions for sturgeon, which were presumably similar in these reaches before regulation and permitted unimpeded movements of sturgeon, thenceforth differed greatly above and below the dam.

Methods

Shovelnose sturgeon were captured from the Williston–Yellowstone and Bismarck segments by using gill nets and trammel nets during the spring and summer. Gill nets were 30.5 m long by 1.8 m deep, with 7.6 cm mesh. Trammel nets were 45.7 m long 3 1.8 m deep, with 5.1-cm inner and 30.5-cm outer mesh. In addition, fish were sampled from the Williston–Yellowstone segment during spring creel censuses. A total of 736 shovelnose sturgeon were sampled from the two segments: 580 fish from the Williston–Yellowstone segment and 156 fish from the Bismarck segment. Fish were sampled from the Williston–Yellowstone segment in 1991, 1992, 1993, 1995, and 1996 and from the Bismarck segment in 1995 (Table 1). Each shovelnose sturgeon was measured for fork length and weighed. Gender was determined for most of the creel fish.

The first ray of the left pectoral fin was removed for age analysis. Collins and Smith (1996) reported that the complete removal of the first pectoral fin ray was nondeleterious in both shortnose sturgeon *Acipenser brevirostrum* and Atlantic sturgeon *A. oxyrinchus*. Our method was less invasive in that we removed only a portion of the pectoral fin ray. The rays were clipped at the point of attachment and again distally about 2.5 cm from this point.

TABLE 1.—Summary of the number of pectoral fin rays collected, aged, and removed from analysis because of age discrepancies.

Year	Collected	Removed	Aged
Williston–Yellowstone			
1991	104	2	
1992	241	10	
1993	87	1	
1995	53	0	
1996	95		




FIGURE 2.—Age-frequency distributions for (A) creel-sampled shovelnose sturgeon from the Williston–Yellowstone segment, (B) netted shovelnose sturgeon from the Williston–Yellowstone segment, and (C) netted shovelnose sturgeon from the Bismarck segment.

ratio, creeled fish within the Williston–Yellowstone were not combined with netted fish within that segment when comparing growth rate differences between segments. Therefore, the subsequent analysis was based on 303 fish of unknown sex netted from the Williston–Yellowstone segment and 145 fish of unknown sex netted from the Bismarck segment.

For netted fish, shovelnose sturgeon in the Williston–Yellowstone segment grew significantly faster than fish in the Bismarck segment (Figure 5). Therefore, the null hypothesis that shovelnose sturgeon had equal growth rates between segments was rejected. The likelihood ratio test for difference between river segments had a test statistic value of 638.8 on 3 degrees of freedom, yielding




FIGURE 3.—Length-frequency distributions for (A) creel-sampled shovelnose sturgeon from the Williston–Yellowstone segment, (B) netted shovelnose sturgeon from the Williston–Yellowstone segment, and (C) netted shovelnose sturgeon from the Bismarck segment.

a *P*-value far less than 0.0001. The fork length of an age-10 fish in the Williston–Yellowstone segment averaged 689 mm, compared with only 457 mm in the Bismarck segment. After age 10, differences in growth rate between the two segments were less pronounced, but fish in the Bismarck segment remained smaller than in the Williston–Yellowstone segment.

Discussion

The mean lengths at age of shovelnose sturgeon vary greatly with geographical location (Figure 6). The mean lengths at age of shovelnose sturgeon from the slightly altered Williston–Yellowstone segment are similar to those reported for shovelnose sturgeon in the Missouri River, Montana, above the confluence of the Yellowstone River




FIGURE 4.—Von Bertalanffy growth (VBG) curves of male and female shovelnose sturgeon from the Williston–Yellowstone segment.

(Gardner and Stewart 1987). Similarly, the mean lengths at age of shovelnose sturgeon from the highly altered Bismarck segment are similar to those reported for shovelnose sturgeon in the Missouri River, South Dakota, below Gavins Point Dam, Nebraska (Zweiacker 1967).

Our length distribution results are consistent with those of Elstad et al. (1992), who reported that shovelnose sturgeon netted in the Williston–Yellowstone segment attained a greater length (maximum FL, 910 mm) than the largest fish netted in the Bismarck segment (maximum FL, 710 mm).




FIGURE 5.—Von Bertalanffy growth curves of shovelnose sturgeon from the Williston–Yellowstone and Bismarck segments.




FIGURE 6.—Mean lengths at age for different shovelnose sturgeon populations in the Missouri and Mississippi rivers: 1, Missouri River above Fort Peck Reservoir, Montana (from Berg 1981); 2, Missouri River above the confluence with the Yellowstone River, Montana (from Gardner and Stewart 1987); 3, Williston–Yellowstone segment of the Missouri and Yellowstone rivers; 4, Bismarck segment of the Missouri River; 5, Missouri River above Vermillion, South Dakota (from Zweigacker 1967); and 6, Pool 13 of the Mississippi River, Iowa (from Helms 1974).

They found that the length distribution in the Williston–Yellowstone segment (range, 320–910 mm FL) also included smaller, presumably younger, fish (320–500 mm), whereas these fish were absent in the narrower length distribution (500–710 mm) of the Bismarck segment. They attributed the narrower distribution in the Bismarck segment to slower growth and the inadequate reproduction and recruitment of small fish. Our results indicated that although fish in the Bismarck segment were shorter than in the Williston–Yellowstone segment, they were not younger but simply grew more slowly (Figure 3).

Habitat changes in the Bismarck segment may explain much of the difference in growth observed between the two segments. After the closure of Garrison Dam in December 1953, the Bismarck segment underwent major changes in water temperature (Figure 7A), hydrograph (Figure 7B), and water clarity (Figure 7C). Hypolimnetic releases from Lake Sakakawea resulted in lower temperatures in the Bismarck segment. As of 1996, mean monthly water temperature in the Missouri River at Bismarck during the peak growing season (June–August) averaged about 78°C colder (148°C

versus 218°C) than in the Yellowstone River at Sidney, Montana. The effect of temperature on fish growth is well documented. Gershanovich (1983) observed a twofold increase in metabolism for sheepshead sturgeon *A. nudipectus* over a temperature increase of 12°C (from 12°C to 24°C). Cech et al. (1984) showed that growth of age-1 white sturgeon was greater at 20°C than at 15°C. Lutes et al. (1990) showed that growth was greater at 18.4°C than at 14.7°C for white sturgeon larvae. Furthermore, Fortin et al. (1996) showed a positive relationship between mean annual air temperature and total length in different populations of lake sturgeon. An increase of 18°C resulted in an average increase of 40 mm for a lake sturgeon between ages 23 and 27. In addition, Nilo et al. (1997) found significant ($P < 0.05$) positive correlation between warmer water and year-class strength in lake sturgeon.




FIGURE 7.—(A)

directly through effects on food supply. Irvine (1985) showed that artificial flow perturbations decreased benthic invertebrate densities. Increased channelization and reduction in backwater areas have led to decreased benthic invertebrate densities in the Missouri River (Hesse and Mestl 1993). During periods of low water, invertebrates concentrate in backwater sloughs in high concentrations. Modde and Schmulbach (1977), studying shovelnose sturgeon from the Missouri River below Gavins Point Dam, Nebraska, suggested that increased water velocities mobilize shovelnose sturgeon prey species, thereby resulting in decreased concentration of food. Furthermore, they found shovelnose sturgeon had lower condition factors during periods of high discharge. In contrast, Megargle and White (1997) found no evidence of discharge affecting shovelnose sturgeon prey items above Fort Peck Reservoir, Montana. Their study area had a natural hydrograph with a relatively large shovelnose sturgeon population that grew to a large size (as much as 3.7 kg).

The lower growth rates observed in the Bismarck segment could have several effects on the population. Slow growth increases the time smaller sturgeon are vulnerable to predation. In addition, marked delays in maturation may occur, especially in a late-maturing species such as shovelnose sturgeon (Hurley and Nickum 1984). Slower growth results in lower production from the population. Moreover, the lack of young shovelnose sturgeon (year-classes 1–4) in the samples collected from the Bismarck segment suggests low recruitment and reproduction. In contrast, the habitat conditions in the Williston-Yellowstone segment serve as a good model for shovelnose sturgeon management. Shovelnose sturgeon from this segment showed faster growth rates and an abundance of diverse year-classes and supported a successful recreational fishery in Montana. Reproduction and recruitment of this population is also occurring; age-1 and age-2 fish have been incidentally caught at Intake, Montana, by paddlefish *Polyodon spathula* snaggers every year since the early 1990s (Scarnecchia, unpublished).

Further information is needed to clarify the effects of habitat alteration on shovelnose sturgeon growth. Techniques need to be refined to capture age-0 and age-1 shovelnose sturgeon more effectively. Specific dietary differences of sturgeon between the two segments of river environment need to be described. In addition, differences in prey selectivity as well as prey availability need to be

identified in these segments as well as throughout the Missouri River. Finally, the relationship between habitat characteristics and forage abundance needs to be determined. Information in these areas will help to clarify the mechanism through which shovelnose sturgeon growth differs between the two river segments.

Acknowledgments

We thank B. Jarrett for his laboratory assistance. This study was supported by the North Dakota Game and Fish Department.

References

- Berg, R. K. 1981. Fish populations of the wild and scenic Missouri River, Montana. Montana Department of Fish, Wildlife and Parks, Federal Aid to Fish and Wildlife Restoration, Project FW-3-R, Job I-A, Helena.
- BioSonics. 1987. Optical pattern recognition system (OPRS), data acquisition program, version 1.08. BioSonics, Seattle.
- Brennan, J. S., and G. M. Cailliet. 1989. Comparative age-determination techniques for white sturgeon in California. Transactions of the American Fisheries Society 118:296–310.
- Brennan, J. S., and G. M. Cailliet. 1991. Age-determination and validation studies of white sturgeon, *Acipenser transmontanus*, in California. Pages 209–234 in P. Willott, editor. *Acipenser*. Centre Tj ect-6-6--

- for floodplain management on the lower Missouri River. *Fisheries* 20(6):30.
- Gardner, W. M., and P. A. Stewart. 1987. The fishery of the Lower Missouri River, Montana. Montana Department of Fish, Wildlife and Parks, Federal Aid to Fish and Wildlife Restoration, Project F-46-R-5, Study 3, Helena.
- Gershanovich, A. D. 1983. Effects of temperature on metabolism, growth and food requirement of young beluga, *Huso huso*, and sheap sturgeon, *Acipenser nudiiventris* (Acipenseridae). *Soviet Journal of Ecology* 21:216–222.
- Helms, D. 1974. Shovelnose sturgeon in the upper Mississippi River, Iowa. Iowa Conservation Commission, Research Technical Series 74-3, Des Moines.
- Hendrickson, J. C., and J. L. Lee. 2000. Aquatic investigations of the Missouri mainstem in North Dakota. North Dakota Game and Fish Department, Fisheries Investigations Report 36, Bismarck.
- Hesse, L. W. 1987. Taming the wild Missouri River: what has it cost? *Fisheries* 12(2):2–9.
- Hesse, L. W., and G. E. Mestl. 1993. An alternative hydrograph for the Missouri River based on the pre-control condition. *North American Journal of Fisheries Management* 13:360–366.
- Hesse, L. W., J. C. Schmulbach, J. M. Carr, K. D. Keenlyne, D. G. Unkenholz, J. W. Robinson, and G. E. Mestl. 1989. Missouri River fishery resources in relation to past, present, and future stresses. Canadian Special Publication of Fisheries and Aquatic Sciences 106:352–371.
- Holton, G. D. 1990. A field guide to Montana fishes. Montana Department of Fish, Wildlife and Parks, Helena.
- Hoopes, D. 1960. Utilization of mayflies and caddisflies by some Mississippi River fish. *Transactions of the American Fisheries Society* 89:32–34.
- Hurley, S. T., and J. G. Nickum. 1984. Spawning and early life history of shovelnose sturgeon. Iowa Co-operative Fish and Wildlife Research Unit, Iowa State University, Project Segment 2-399-R-1, Completion Report, Ames.
- Hyndes, G. A., and I. C. Potter. 1996. Comparison between the age structures, growth and reproductive biology of two co-occurring sillaginids, *Sillago robusta* and *S. bassensis*, in temperate coastal waters of Australia. *Journal of Fish Biology* 49:14–32.
- Irvine, J. R. 1985. Effects of successive flow perturbations on stream invertebrates. *Canadian Journal of Fisheries and Aquatic Sciences* 42:1922–1927.
- Kimura, D. K. 1980. Likelihood methods for the von Bertalanffy growth curve. *U.S. National Marine Fisheries Service Fishery Bulletin* 77:765–776.
- LeBreton, G. T. O., and F. W. H. Beamish. 2000. Suitability of lake sturgeon growth rings in chronological development. *Transactions of the American Fisheries Society* 129:1018–1030.
- Lee, D. S. 1978. *Scaphirhynchus platyrhynchus* (Rafinesques). Page 44 in D. S. Lee, C. R. Gilbert, C. H. Hocutt, R. E. Jenkins, D. E. McAllister, and J. R. Stauffer, Jr., editors. *Atlas of North American freshwater fishes*. North Carolina State Museum of Natural History, Raleigh.
- Lutes, P. B., S. S. O. Hung, and F. S. Conte. 1990. Survival, growth, and body condition of white sturgeon larvae fed purified and commercial diets at 14.7 and 18.4 degrees Celsius. *Progressive Fish-Culturist* 52:192–196.
- Megargle, D. J., and R. G. White. 1997. Temporal variation in diet and food selection of shovelnose sturgeon in the Missouri River above Fort Peck Reservoir, Montana. Montana State University Completion report to U.S. Army Corps of Engineers, Omaha District, Omaha, Nebraska.
- Modde, T. C. 1971. Food selectivity of the shovelnose sturgeon *Scaphirhynchus platyrhynchus*, in the Missouri River. Master's thesis. University of South Dakota, Vermillion.
- Modde, T. C., and J. C. Schmulbach. 1977. Food and feeding behavior of the shovelnose sturgeon, *Scaphirhynchus platyrhynchus*, in the unchannelized Missouri River, South Dakota. *Transactions of the American Fisheries Society* 106:602–608.
- Nilo, P., P. Dumont, and R. Fortin. 1997. Climatic and hydrological determinants of year-class strength of St. Lawrence River lake sturgeon (*Acipenser fulvescens*). *Canadian Journal of Fisheries and Aquatic Sciences* 54:774–780.
- Pflieger, W. L. 1975. The fishes of Missouri. Missouri Department of Conservation, Jefferson City.
- Rossiter, A., D. L. G. Noakes, and F. W. H. Beamish. 1995. Validation of age estimation for the lake sturgeon. *Transactions of the American Fisheries Society* 124:777–781.
- SAS Institute. 1989. SAS/OR user's guide, version 6. SAS Institute, Cary, North Carolina.

- J. R. Zuboy, editors. Proceedings of the symposium on restoration planning for rivers of the Mississippi River ecosystem. National Biological Survey, Biological Report 19, Washington, D.C.
- Zweiacker, P. L. 1967. Aspects of the life history of the shovelnose sturgeon, *Scaphirhynchus platyrhynchus* (Rafinesque) in the Missouri River. Master's thesis. University of South Dakota, Vermillion.