

Figure 3

www.elsevier.com/locate/jmb

choof on February 3, 1992.

the
l
a
lie
in
t
oo
h
of

ed p
es l
i
w
f r

w
ne
h
y
s
ar
ea

conducted at
in 1965 and
in 1968. Their
conclusions
from 1964 were
that the in-
creased use
of organic and
inorganic

with a 1990 census
of the 13,000 com-
munities in the United States, the
percentage of the population that
lived in urban areas rose from 49.5
percent in 1980 to 53.5 percent in
1990. The percentage of the popu-
lation that lived in rural areas fell
from 49.5 percent in 1980 to 46.5
percent in 1990. The percentage of
the population that lived in sub-
urban areas rose from 1.0 percent
in 1980 to 1.5 percent in 1990.

1. **Point sampling** is a technique that involves randomly selecting points within a study area and then recording the number of individuals of a particular species at each point. This method is useful for estimating the density of a population in a large area.

to find a way to make the
diesel engine and play
right through the oc-
casion conditions
to and be
the old

946. A sur-
titio e. h. of
o. h. of
factar

3-1989

ep
 19%
 Agricultural, B

Is the child of a
single parent better

City, 2000

[illegible]

ine Valley, based on
with a some risk
leaf blade b. k. + 15
of end, 100% of 10
age did not increase
white leaf blade tiss
content, boron, shoul

the best field trials
the, including both
1994-70
beet fields in th
including both

the nutrient status of
terrestrial biotopes

Chemistry and Soil

Office of applied horticulture
Station, Research

[illegible][illegible]




Figure 1. Sugar % as affected by preplant applied boron




Figure 2. Conductivity as affected by preplant applied boron.