,^												
1				胡桃	指旗			34				illigations' en part
į į		作监查		黄疸	ily In							
,			鸖		11	::밥.				11		(1일부) - 1일(조)
D A IDI			ú /l	erang. Pada sa								
DAIRY	COMPOS			19 79 19 14	s in the following of t					, j		
`*		k (176)	di [中长	Jyr	ជ្រាប់			pi Kaji			
1, 7, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,					基式		l In S				ķi j	$\begin{bmatrix} 3 & 1 \end{bmatrix}$
Onivers.	y of Idaho 2Wilbur-E	*		序数	Wid				루푸	الإلا		
4	Wilder-E. his Cooking still a			用增	\W	: {\pi			Said Maria d			
								1 11				3.177
osted dairy	Sanura Wa					1		i ja	 	불다	Mr. And	sativa
- 1	64 - 1966 - 1968 1968				1	; di	i ili	3 \ 2 1	ngi i			oluble
lized plots v	mpact on all fall fall fall fall fall fall fall			1				ا مولد اد		[i]	Jay / Science	Dmpost
phosphate fe	11170r con				ll) It'						ri Valdoriya	of soil,
ch of the P, p	「Y · · · · · · · · · · · · · · · · · · ·		, liki						強し	智贷点		can be
compost.	massium (150), and control e		游员		盟鄉	F	[11][일 D 호] [1] [1] 보		jeir.			i can oc
Joinpost.		歷史			Ģď,) / (a)		
N.			y H	$h_{ij}(i)$	13				苏萨	i (III 🕌	White!	Tit
ost is an en	ronmental							国教的 医别维	學工學		Michigan.	ical to
nutrients fr	[2]				ŊĠ						3. 其其第二	er than
	[6] ····		h in			HrMT:		\mathbb{R}^{N}		為農业	, coenty áir sei	better
y applying ς	mpost to		[발발] 교 보호		ار این			i_s				ported
y forage har	i sting.		ji †¹		ang sport Til state Til state			g Vill		并靠		1000 PM. *
osting man	e can reduce field test	1 1		i is ill oradii	. 			il.				nanure
y reducing	rolume and undistricted a		11 (1 11 (4)		t nei						s zaposaju. Esabbado	crease
niformity du	to a reduc				1 4	1 (No. 14) 100 (No. 14)		$ P_0\rangle$:	viable
nd phytotox	Substance of the substa			[115] 						. Úd.		there
y greater pr	duction ar a construction are		1. 1.		1 1		the feet		$\frac{1}{4}$	型 (制) - 25 jii		ig and
s of nutrient				elija. Politika		Ţ.,						
bjectives of	ne study w		1 1		$\int_{0}^{\infty} \left\{ -\frac{1}{2} \right\}$	i li		B.			. 100 9a da - 200 con	rtneuf
with alfalfa	n southerr			顶升								des to
5 distalliable				131 		Paka i			有特	k		ycling
n dairy man		1.16		$J_{ij}^{(i)}(\vec{s}, \vec{s})$			Hill		例。	10.6		
بالن				ji' [闡達				17 1	ledică. Nation		
	ja – Principalis kalturian karandar kalturi	ч : <mark>1</mark>	加拉	集集	4				h.	, f		
<i>on</i> Ultra' varie⊯	of alfalf					ÿ ∦İ			1-		数据证	ıfter a
of dry edit:	hears (Particular and a second		6. I		1, 1					i211	12001	iberly,
oil was a sur		F 31 2.1	1-11		贈		HE			机溶剂	Laker	inodic
alcid) near I	Fig. 1. Superior Control of the property of the control of the	門門		蓝绿	الزوار و مرج	T. Accellant					Mich II	. The
design was	[2] Tak. Allela L. Mar. (1996). The decidal of the Park No. (1996).				ր յ (撰				\$17 7 73 { 53711		
						31	l l _e i ety li		見りま			
18 10 10		1 i () 			24 (4 (14.14)				5 J.		. · · ·	
ted treatmen	were: an		1.1	2017 2017	: 1년 10 년							e (5.6
y compost p	or to estab				(19)				jidh	 	a Bugali a	er acre
1) compost 11	- 4 1 29/2 E 1 11 38 2002 1 1 1 1 1 1		Ji da		' -						, albertaan 98. See Geekneemee	innual
f 2.5 tons i	ir acre (5. 10. 10. 10. 10. 10. 10. 10. 10. 10. 10	1									jim udejroesiako Lijudika santolia	er (F)
ovide nutric		100			١	4/1				i i	E POR LEGIS LEGISTORIA	' Hor to
and annuall			$ \cdot _{\mathbb{T}_q}$		i qar				į, į,	두다	- 10 000 - 10 000	g ha ⁻¹)
) ()	Make the model from the first of the track of the first o	[1,1]	1, 5	· .	1	11	1		1. 7.1	1.	They wanter	[define]
į.		∏ tat					get i			1		78
uji S	4	in 1					, 1		[]		J j	/0
	· · · · · · · · · · · · · · · · · · ·		事事				proje			1		
· .			<u>, , , </u>	<u> </u>	<u></u>	<u> </u>	: []. <u>- </u>		a (<u>, </u>	

monoammonium phosphate (11-52-0). Annual treatments were repeated every year in the same plot so that cumulative treatment effects on soil test nutrients and yield over 3 years could be determined.

Composted dairly manure was applied to the soil prior to planting at a standard rate of compost (C1X) and twice the standard rate (C2X), and incorporated to a 6 inch (15-cm) depth prior to seeding alfalfal Different batches of compost were applied in subsequent years to the same plots (4 x 2 m) at 0 (control), 4 for C1X, and 8 tons per acre for C2X treatments (8.93 and 17.9 Mg DM ha⁻¹ yr⁻¹, respectively) on 10 April 2001. In 2002 the actual rates were 2.22 (C1X) and 4.44 tons per acre (C2X) (5.0 and 110.0 Mg ha⁻¹) on an as-delivered basis.

A 30 by 20-foot (9.1 by 6.1-n) to rp was used to collect and weigh compost spread by the truck to determine actual application rates for each batch. A commercial compost spreader truck applied the compost in andomized 3 3-fd ot (9.1-m) wide strips, and plots were established within the strips as 30 by 100-10ot (9.1 by 30.4-m) plots.

Soil and Compost Sampling and Analys es

Soil samples were taken from the field using a Giddings Probe and a 1.25 inch (3-cm) hollow core sampler. Samples were taken in October 2000, before treatment initiation; March of 2001, 2002, and 2003; and October 2000? and 2003 at the conclusion of the study. Each time, 7 soil cores were composited for each plot from 0 to 3, 3 to 12, and 12 to 24 inches (0 to 7.6, 7.6 to 30.5, and 30.5 to 61 cm). Samples were analyzed for Olsen P.

Compost samples were collected as grab samples from the tarp and composited for each batch, and stored frozen until analyzed.

RESUL'IS

Compost Characteristics

The composition of the compost was similar in both 2001 and 2002. The average nutrient analysis and application of nutrients for the dairy compost is shown in Table 1.